

Communication

**Tetranuclear Manganese Models of the OEC Displaying  
Hydrogen Bonding Interactions: Application to  
Electrocatalytic Water Oxidation to Hydrogen Peroxide**

Zhiji Han, Kyle T. Horak, Heui Beom Lee, and Theodor Agapie

*J. Am. Chem. Soc.*, **Just Accepted Manuscript** • Publication Date (Web): 06 Jun 2017

Downloaded from <http://pubs.acs.org> on June 8, 2017

**Just Accepted**

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



**ACS Publications**

# Tetranuclear Manganese Models of the OEC Displaying Hydrogen Bonding Interactions: Application to Electrocatalytic Water Oxidation to Hydrogen Peroxide


Zhiji Han, Kyle T. Horak, Heui Beom Lee and Theodor Agapie\*

Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, United States

## Supporting Information Placeholder

**ABSTRACT:** Toward the development of structural and functional models of the oxygen evolving complex (OEC) of photosystem II (PSII), we report the synthesis of site-differentiated tetranuclear manganese complexes featuring three six-coordinate and one five-coordinate Mn centers. To incorporate biologically relevant second coordination sphere interactions, substituents capable of hydrogen bonding are included as pyrazolates with phenylamino groups. Complexes with terminal anionic ligands,  $\text{OH}^-$  or  $\text{Cl}^-$ , bound to the lower coordinate metal center are supported through the hydrogen-bonding network in a fashion reminiscent to the enzymatic active site. The hydroxide complex was found to be a competent electrocatalyst for O-O bond formation, a key transformation pertinent to the OEC. In an acetonitrile-water mixture, at neutral pH, electrochemical water oxidation to hydrogen peroxide was observed, albeit with low (15%) Faradaic yield, likely due to competing reactions with organics. In agreement, 9,10-dihydroanthracene is electrochemically oxidized in the presence of this cluster both via H-atom abstraction and oxygenation at ~50% combined Faradaic yield.


Water oxidation as source of reducing equivalents for artificial photosynthesis is an appealing approach to solar energy conversion and storage.<sup>1</sup> The oxygen evolving complex (OEC) of photosystem II (PSII) is the biological catalyst for water oxidation to  $\text{O}_2$ .<sup>2</sup> The OEC features a multinuclear active site,  $\text{CaMn}_4\text{O}_5$ , consisting of a cubane  $\text{CaMn}_3\text{O}_4$  moiety connected to a “dangling” Mn center; this cluster coordinates water ligands that are engaged in hydrogen bonding networks (Figure 1A).<sup>2a, 3</sup> Significant efforts have been focused on elucidating the mechanism of water oxidation by the OEC in PSII.<sup>2a, 4</sup> Given the complexity of PSII and the challenge to perform structure-function studies with the biological system, discrete cluster models have been targeted to provide insight into the function of the OEC.<sup>5</sup> Synthetic cluster models that display both Ca and Mn centers and a cubane moiety have been prepared, though they do not display catalytic activity.<sup>6</sup> Multinuclear Mn complexes capable of O-O bond formation are rare.<sup>2b, 2d, 7</sup> More generally, several discrete clusters of first row transition metals with structures reminiscent of the OEC have been reported to support  $\text{O}_2$  evolution, although *in situ* formed heterogeneous metal oxides can be responsible for catalysis.<sup>2b, 7g-i, 8</sup> In some cases, the substrate oxygen has been proposed to be a terminal metal-oxo species.<sup>2a, 2b</sup> Deprotonations are an integral part of the conversion of water to such moiety. Furthermore, challenging redox steps can be facilitated by coupling of electron and proton transfers.<sup>9</sup> Toward gaining insight into the roles of



**Figure 1. A)** Drawing of the OEC (PDB 4UB8) highlighting a tetranuclear cluster including the dangling Mn (orange) and H-bonding network (blue)<sup>3b,d</sup>; **B)** Target synthetic model.

protons in the chemistry of transition metals with oxygen, a variety of elegant strategies have been reported for supporting transition metal oxo and hydroxo chemistry with hydrogen bonding networks in mononuclear complexes.<sup>10</sup> Such strategies are significantly limited in the context of multinuclear complexes. Moreover, terminal oxo or hydroxo ligands are rare in multinuclear models of the OEC.<sup>11</sup> Herein, we report a synthetic strategy to access a  $\text{Mn}_4$  cluster model of the OEC that has a dangling Mn motif, a terminal OH ligand, and second coordination sphere H-bonding to stabilize it (Figure 1B). This species demonstrates O-O bond formation under electrochemical conditions.

Tetranuclear  $\text{Mn}_4$  complexes were prepared in a one pot procedure starting from trinuclear precursor  $\text{LMn}_3(\text{OAc})_3$  **1** (Scheme 1). Addition of  $\text{Ca}(\text{OTf})_2$  (1.6 equiv) and  $[\text{RNHPz}]\text{Na}$  (3.5 equiv) to **1**,<sup>12</sup> followed by iodosobenzene (PhIO) (1.0 equiv) and  $\text{Mn}(\text{II})$  salt (1-2 equiv) results in the desired tetramanganese complexes.  $\text{Ca}(\text{OTf})_2$  facilitates the substitution of acetate ligands with pyrazolates. PhIO acts as the O-atom source for the incorporation of the interstitial oxide moiety. The additional equivalent of  $\text{Mn}(\text{II})$  is the source of apical metal in the final cluster.<sup>13</sup> Using  $[\text{PhNHPz}]\text{Na}$  as the bidentate ligand and  $\text{MnCl}_2$  as  $\text{Mn}(\text{II})$  precursor gave  $[\text{LMn}_3(\text{PhNHPz})_3\text{OMnCl}][\text{OTf}]$  (**2**) in ~55% isolated yield. ESI-MS and single crystal X-ray diffraction (XRD, *vide infra*) characterization is consistent with this assignment. A similar synthetic strategy was applied to target a lower coordinate or triflate complex by using  $\text{Mn}(\text{OTf})_2$  as the fourth equivalent source. ESI-MS characterization of this product suggests the formation of a hydroxide complex. X-ray diffraction quality crystals of this compound have not been obtained to date using the  $[\text{PhNHPz}]\text{Na}$  as the bridging ligand. However, using  $[4\text{-tBuC}_6\text{H}_4\text{NHPz}]\text{Na}$  resulted in a complex with ESI-MS data corresponding to a hydroxide species ( $[\text{LMn}_3(4\text{-tBuC}_6\text{H}_4\text{NHPz})_3\text{OMn}(\text{OH})][\text{OTf}]$ , **3**) that gave crystals suitable for XRD studies.




**Scheme 1.** Synthesis of tetranuclear manganese complexes and crystal structures of **2** and **3**. Thermal ellipsoids are shown at the 50% probability level. Parts of the ligands, hydrogen atoms, triflate, and co-crystallized solvent molecules are not shown for clarity.

Solid-state structures of **2** and **3** were determined by XRD (Scheme 1). In both structures, the three basal Mn centers are in a distorted octahedral coordination environment with two bridging alkoxides, one pyrazolate, two pyridine, and a  $\mu_4$ -oxo ligands each. The apical Mn displays a trigonal bipyramidal geometry and is connected to the three basal Mn centers through the interstitial  $\mu_4$ -oxo ion. The Cl(1)-N(1) distance (3.056(7) Å) in **2** is slightly shorter than the range (3.238–3.326 Å) observed in a previously reported five coordinate mononuclear Mn-Cl complex with intramolecular hydrogen bonding networks indicating that all NH bonds engage in dative hydrogen bonding interactions with the chloride.<sup>14</sup>

The Mn- $\mu_4$ -oxo distances in **3** are consistent with the presence of two Mn(II) (Mn(3) and Mn(4)) and two Mn(III) (Mn(1) and Mn(2)).<sup>13</sup> These oxidation state assignments, charge balance, and localization of OH and NH hydrogens in the electron density map support the assignment of the apical ligand as a hydroxide. Compound **3** displays two short O-N distances between the hydroxide oxygen and the aniline nitrogens (2.774(5) and 2.805(7) Å, for N(1) and N(2), respectively). These distances correspond to strong hydrogen bonding interactions, in the range previously reported (2.721–2.848 Å) for mononuclear Mn(III)-oxo species.<sup>10a</sup> The geometry at nitrogen is consistent with the two anilines acting as hydrogen bond donors to the hydroxide lone pairs. The third O-N distance is longer (2.939(5) Å, to N(3)), and the orientation is consistent with the aniline serving as the hydrogen bond acceptor. The N(3)H bond acts as a hydrogen bond donor to a triflate ion. The differences in the hydrogen bonding interactions in **2** vs **3** highlight the versatility of the aniline moiety in accommodating both donor and acceptor functions. The Mn(1)-O(2) distance (1.874(3) Å) is comparable to a mononuclear complex displaying an intramolecular hydrogen bonding network, at 1.872(2) Å, suggesting that the coordination environment for the Mn(1)-OH moiety is similar to monomanganese species despite being part of a cluster.<sup>10a</sup> All structural parameters are consistent with a cluster supporting a Mn(III)-coordinate terminal hydroxide that is engaged in several hydrogen bonds. Although the geometry of the donors and acceptors

differs, the structural parameters are similar between the OEC and **3**; the three non-hydrogen atoms closest to one of the OEC terminal oxygenic ligand are in the range of 2.67–3.04 Å (Figure 1),<sup>3d</sup> which includes the range observed in **3** for the O(2)-N distances.

The electrochemical properties of **3** were investigated by cyclic voltammetry (Figure 2). Two reversible one-electron redox events are observed at -0.72 and -0.21 and a quasireversible redox couple at 0.35 V vs Fc/Fc<sup>+</sup>. In the context of water oxidation by the OEC, the electro-



**Figure 2.** Cyclic voltammograms of 1.0 mM of [LMn<sub>3</sub>(4-*t*BuC<sub>6</sub>H<sub>4</sub>NHPz)<sub>3</sub>OMn(OH)][OTf] (**3**) in MeCN without water (gray dash) and in the presence of water with increasing concentration (other colors). Conditions: 0.1 M [*n*-Bu<sub>4</sub>N][ClO<sub>4</sub>], GC working electrode, 0.07 cm<sup>2</sup>, Pt wire counter electrode, scan rate: 100 mV/s.

chemical behavior of **3** in the presence of water is of interest. Independently, compound **3** was found to be soluble and stable in MeCN/H<sub>2</sub>O mixtures containing up to 10 % volume water as observed by <sup>1</sup>H NMR spectroscopy (Figure S13). The CV of **3** was recorded in the presence of various concentrations of water (Figure 2). Increased concentration of water correlates with increased current close to the second oxidation event, which is assigned to a catalytic process.

The catalytic wave observed upon addition of water suggests that compound **3** acts as water oxidation catalyst. In order to quantify the reaction products upon electrochemical oxidation, we performed controlled potential electrolysis (CPE) experiments at 0.57 V (vs Fc/Fc<sup>+</sup>). Complex **3** was dissolved in a mixture of MeCN/H<sub>2</sub>O (19:1 v/v) with 0.1 M [*n*-Bu<sub>4</sub>N][ClO<sub>4</sub>] as supporting electrolyte. The experiments were performed in a two-compartment cell separated by an anion-exchange membrane. In a typical experiment, with a total amount of 5.8 C passed charge over the course of 1.5 h (Figure S9), H<sub>2</sub>O<sub>2</sub> was detected with a Faradaic efficiency of ~15% using a chemiluminescence method with 10-acetyl-3,7-dihydroxyphenoxazine in combination of horseradish peroxidase (Figure S1).<sup>15</sup> Addition of catalase to the reaction mixture after electrolysis produced O<sub>2</sub> as detected by gas chromatography, which is also consistent with the generation of H<sub>2</sub>O<sub>2</sub> by **3** (Figure S2). The exact overpotential of the system remains undetermined due to the use of a mixture of solvents. The potential for water oxidation to O<sub>2</sub> drifts ~ 200 mV more positive in acetonitrile,<sup>16</sup> for example.

The low observed Faradaic yield for H<sub>2</sub>O<sub>2</sub> formation could be due to side reactions between the highly reactive species generated upon oxidation, such as Mn-oxo or Mn-(hydro)peroxo species, with organic moieties. To test this hypothesis, CPE was conducted in the presence of 9,10-dihydroanthracene (0.1 M) as a source of weak C-H bond to react with the intermediates. H<sub>2</sub>O<sub>2</sub> was not detected in this experiment and the combined Faradaic efficiency corresponding to 9,10-dihydroanthracene oxidation to anthracene and anthraquinone was 51% (Figure S10). CV control experiments show that there is no oxidative wave for 9,10-dihydroanthracene in the absence of **3** at the applied potential of CPE experiments (Figure S11). Stirring 9,10-dihydroanthracene in the presence of **3** without applied potential also results in no oxidized products. These observations are consistent with the generation of reactive intermediates that perform H-atom abstraction and oxygenation reactions of C-H bonds competitive with water oxidation. The lack of H<sub>2</sub>O<sub>2</sub> formation and the significant increase of the Faradaic yield in the presence of 9,10-dihydroanthracene indicate a faster reaction between the intermediates and the weak C-H bonds of this substrate than both H<sub>2</sub>O<sub>2</sub> formation and side reactions with the organic solvent.

To evaluate the stability of **3** during catalysis in more detail, electrochemical experiments reveal that the voltammogram traces remain unchanged for 200 continuous cycles over the course of 3 h (Figure S12). After these scans, rinsing the electrode gently without polishing gave no catalytic wave and no redox features in a fresh electrolyte solution (Figure S12). Similar rinsing experiments were performed after CPE and showed no consumption of charge (Figure S9). These observations suggest that there is no deposition of catalytically active metal species on the electrode. Furthermore, to test the possibility of formation of heterogeneous MnO<sub>x</sub> catalysts via free Mn ions, chelating ligands were added to solutions of **3** during CV measurements. No decrease of catalytic current was observed (Figures S14-16).<sup>17</sup> Additionally, dynamic light scattering measurements do not detect any nanoparticles (Figure S20). These observations are consistent with the absence of MnO<sub>x</sub> formation from **3** in our electrochemical experiments. To evaluate whether other clusters that may form under the

reaction conditions for the synthesis of **3** in small concentrations could be the active catalysts, several preparations were tested leaving some of the reagents out (Figures S18-19). In all cases tested, smaller catalytic current was observed compared to **3** at the same concentration. Thus, the control experiments are consistent with **3** as the active molecular catalyst for water oxidation to H<sub>2</sub>O<sub>2</sub>.

In summary, we report the synthesis of tetranuclear manganese complexes displaying hydrogen bonding networks as more accurate structural models of the biological water oxidation catalyst that include second coordination sphere interactions. In particular, a cluster displaying a terminal hydroxide ligand is reminiscent of structural motifs involving the dangler Mn center of the OEC. Hydrogen bond donors and acceptors stabilize the terminal hydroxide ligand. This species supports electrocatalytic water oxidation to H<sub>2</sub>O<sub>2</sub>, albeit in low yield, in a rare demonstration of function (O-O bond formation) with a structural model of the OEC. Related to the mechanism of the biological system, the present results suggest that a terminal Mn-hydroxide moiety can serve as precursor to O-O bond formation chemistry. Importantly, control experiments are inconsistent with the generation of a heterogeneous active species, and support complex **3** as a discrete and well-defined electrocatalyst for water oxidation. Complex **3** is also able to perform both H-atom abstraction and oxygenation reactions with 9,10-dihydroanthracene under electrocatalytic conditions. Current studies are focused on the mechanistic aspects of these transformations.

## ASSOCIATED CONTENT

### Supporting Information

This material is available free of charge via the Internet at <http://pubs.acs.org>. Crystallographic data (CIF). General considerations, physical methods, and synthetic procedure. Figures S1-S18, Table S1 (PDF)

## AUTHOR INFORMATION

### Corresponding Author

\*agapie@caltech.edu

### Notes

The authors declare no competing financial interests.

## ACKNOWLEDGMENT

This research was supported by the NIH (R01-GM102687A). We thank L. M. Henling for assistance with crystallography, J.C. Tran for assistance with the characterization of the 4-*t*BuC<sub>6</sub>H<sub>4</sub>NHPzH, and B. M. Hunter for helpful discussion on H<sub>2</sub>O<sub>2</sub> detection.

## REFERENCES

- (a) Lewis, N. S.; Nocera, D. G., *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 15729-15735; (b) Gray, H. B., *Nat. Chem.* **2009**, *1*, 7-7.
- (a) Yano, J.; Yachandra, V., *Chem. Rev.* **2014**, *114*, 4175-4205; (b) Blakemore, J. D.; Crabtree, R. H.; Brudvig, G. W., *Chem. Rev.* **2015**, *115*, 12974-13005; (c) Vinyard, D. J.; Ananyev, G. M.; Dismukes, G. C., *Annu. Rev. Biochem.* **2013**, *82*, 577-606; (d) Kärkäs, M. D.; Åkermark, B., *Dalton Trans.* **2016**, *45*, 14421-14461.
- (a) Ferreira, K. N.; Iverson, T. M.; Maghlaoui, K.; Barber, J.; Iwata, S., *Science* **2004**, *303*, 1831-1838; (b) Umena, Y.; Kawakami, K.; Shen, J. R.; Kamiya, N., *Nature* **2011**, *473*, 55-65; (c) Suga, M.; Akita, F.; Hirata, K.; Ueno, G.; Murakami, H.; Nakajima, Y.; Shimizu, T.; Yamashita, K.; Yamamoto, M.; Ago, H.; Shen, J. R., *Nature* **2015**, *517*, 99-103; (d) Young, I. D.; Ibrahim, M.; Chatterjee, R.; Gul, S.; Fuller, F.



- D.; Koroidov, S.; Brewster, A. S.; Tran, R.; Alonso-Mori, R.; Kroll, T.; al., e., *Nature* **2016**, *540*, 453-457.
4. Cox, N.; Retegan, M.; Neese, F.; Pantazis, D. A.; Boussac, A.; Lubitz, W., *Science* **2014**, *345*, 804-808.
5. Tsui, E. Y.; Kanady, J. S.; Agapie, T., *Inorg. Chem.* **2013**, *52*, 13833-13848.
6. (a) Zhang, C. X.; Chen, C. H.; Dong, H. X.; Shen, J. R.; Dau, H.; Zhao, J. Q., *Science* **2015**, *348*, 690-693; (b) Kanady, J. S.; Lin, P. H.; Carsch, K. M.; Nielsen, R. J.; Takase, M. K.; Goddard, W. A.; Agapie, T., *J. Am. Chem. Soc.* **2014**, *136*, 14373-14376; (c) Mukherjee, S.; Stull, J. A.; Yano, J.; Stamatatos, T. C.; Pringouri, K.; Stich, T. A.; Abboud, K. A.; Britt, R. D.; Yachandra, V. K.; Christou, G., *Proc. Natl. Acad. Sci. U.S.A.* **2012**, *109*, 2257-2262; (d) Kanady, J. S.; Tsui, E. Y.; Day, M. W.; Agapie, T., *Science* **2011**, *333*, 733-736.
7. (a) Limburg, J.; Vrettos, J. S.; Liable-Sands, L. M.; Rheingold, A. L.; Crabtree, R. H.; Brudvig, G. W., *Science* **1999**, *283*, 1524-1527; (b) Naruta, Y.; Sasayama, M.; Sasaki, T., *Angew. Chem. Int. Ed.* **1994**, *33*, 1839-1841; (c) Karlsson, E. A.; Lee, B. L.; Åkermark, T.; Johnston, E. V.; Kärkäs, M. D.; Sun, J. L.; Hansson, O.; Bäckvall, J.; Åkermark, B., *Angew. Chem. Int. Ed.* **2011**, *50*, 11715-11718; (d) Najafpour, M. M.; Renger, G.; Holynska, M.; Moghaddam, A. N.; Aro, E. M.; Carpentier, R.; Nishihara, H.; Eaton-Rye, J. J.; Shen, J. R.; Allakhverdiev, S. I., *Chem. Rev.* **2016**, *116*, 2886-2936; (e) Poulsen, A. K.; Rempel, A.; McKenzie, C. J., *Angew. Chem. Int. Ed.* **2005**, *44*, 6916-6920; (f) Beckmann, K.; Uchtenhagen, H.; Berggren, G.; Anderlund, M. F.; Thapper, A.; Messinger, J.; Styring, S.; Kurz, P., *Energ. Environ. Sci.* **2008**, *1*, 668-676; (g) Ruettinger, W.; Yagi, M.; Wolf, K.; Bernasek, S.; Dismukes, G. C., *J. Am. Chem. Soc.* **2000**, *122*, 10353-10357; (h) Brimblecombe, R.; Kolling, D. R. J.; Bond, A. M.; Dismukes, G. C.; Swiegers, G. F.; Spiccia, L., *Inorg. Chem.* **2009**, *48*, 7269-7279; (i) Dismukes, G. C.; Brimblecombe, R.; Felton, G. A. N.; Pryadun, R. S.; Sheats, J. E.; Spiccia, L.; Swiegers, G. F., *Acc. Chem. Res.* **2009**, *42*, 1935-1943.
8. (a) Evangelisti, F.; More, R.; Hodel, F.; Luber, S.; Patzke, G. R., *J. Am. Chem. Soc.* **2015**, *137*, 11076-11084; (b) Yin, Q. S.; Tan, J. M.; Besson, C.; Geletii, Y. V.; Musaev, D. G.; Kuznetsov, A. E.; Luo, Z.; Hardcastle, K. I.; Hill, C. L., *Science* **2010**, *328*, 342-345; (c) Stracke, J. J.; Finke, R. G., *J. Am. Chem. Soc.* **2011**, *133*, 14872-14875; (d) Brimblecombe, R.; Swiegers, G. F.; Dismukes, G. C.; Spiccia, L., *Angew. Chem. Int. Ed.* **2008**, *47*, 7335-7338; (e) Hocking, R. K.; Brimblecombe, R.; Chang, L. Y.; Singh, A.; Cheah, M. H.; Glover, C.; Casey, W. H.; Spiccia, L., *Nat. Chem.* **2011**, *3*, 461-466; (f) McCool, N. S.; Robinson, D. M.; Sheats, J. E.; Dismukes, G. C., *J. Am. Chem. Soc.* **2011**, *133*, 11446-11449; (g) Berardi, S.; La Ganga, G.; Natali, M.; Bazzan, I.; Puntoriero, F.; Sartorel, A.; Scandola, F.; Campagna, S.; Bonchio, M., *J. Am. Chem. Soc.* **2012**, *134*, 11104-11107; (h) Smith, P. F.; Kaplan, C.; Sheats, J. E.; Robinson, D. M.; McCool, N. S.; Mezle, N.; Dismukes, G. C., *Inorg. Chem.* **2014**, *53*, 2113-2121; (i) Ullman, A. M.; Liu, Y.; Huynh, M.; Bediako, D. K.; Wang, H. S.; Anderson, B. L.; Powers, D. C.; Breen, J. J.; Abruna, H. D.; Nocera, D. G., *J. Am. Chem. Soc.* **2014**, *136*, 17681-17688; (j) Nguyen, A. I.; Ziegler, M. S.; Ona-Burgos, P.; Sturzbecher-Hohne, M.; Kim, W.; Bellone, D. E.; Tilley, T. D., *J. Am. Chem. Soc.* **2015**, *137*, 12865-12872; (k) Smith, P. F.; Hunt, L.; Laursen, A. B.; Sagar, V.; Kaushik, S.; Calvinho, K. U. D.; Marotta, G.; Mosconi, E.; De Angelis, F.; Dismukes, G. C., *J. Am. Chem. Soc.* **2015**, *137*, 15460-15468; (l) Okamura, M.; Kondo, M.; Kuga, R.; Kurashige, Y.; Yanai, T.; Hayami, S.; Praneeth, V. K. K.; Yoshida, M.; Yoneda, K.; Kawata, S.; Masaoka, S., *Nature* **2016**, *530*, 465-468; (m) Cady, C. W.; Crabtree, R. H.; Brudvig, G. W., *Coord. Chem. Rev.* **2008**, *252*, 444-455.
9. Weinberg, D. R.; Gagliardi, C. J.; Hull, J. F.; Murphy, C. F.; Kent, C. A.; Westlake, B. C.; Paul, A.; Ess, D. H.; McCafferty, D. G.; Meyer, T. J., *Chem. Rev.* **2012**, *112*, 4016-4093.
10. (a) Shirin, Z.; Hammes, B. S.; Young, V. G.; Borovik, A. S., *J. Am. Chem. Soc.* **2000**, *122*, 1836-1837; (b) MacBeth, C. E.; Gupta, R.; Mitchell-Koch, K. R.; Young, V. G.; Lushington, G. H.; Thompson, W. H.; Hendrich, M. P.; Borovik, A. S., *J. Am. Chem. Soc.* **2004**, *126*, 2556-2567; (c) Shook, R. L.; Peterson, S. M.; Greaves, J.; Moore, C.; Rheingold, A. L.; Borovik, A. S., *J. Am. Chem. Soc.* **2011**, *133*, 5810-5817; (d) Borovik, A. S., *Acc. Chem. Res.* **2005**, *38*, 54-61; (e) Ford, C. L.; Park, Y. J.; Matson, E. M.; Gordon, Z.; Fout, A. R., *Science* **2016**, *354*, 741-743; (f) Zong, R.; Thummel, R. P., *J. Am. Chem. Soc.* **2005**, *127*, 12802-12803; (g) Dogutan, D. K.; McGuire, R.; Nocera, D. G., *J. Am. Chem. Soc.* **2011**, *133*, 9178-9180; (h) Chang, C. J.; Chng, L. L.; Nocera, D. G., *J. Am. Chem. Soc.* **2003**, *125*, 1866-1876; (i) Carver, C. T.; Matson, B. D.; Mayer, J. M., *J. Am. Chem. Soc.* **2012**, *134*, 5444-5447; (j) Rigsby, M. L.; Wasylenko, D. J.; Pegis, M. L.; Mayer, J. M., *J. Am. Chem. Soc.* **2015**, *137*, 4296-4299.
11. A CSD search of small Mn clusters displaying terminal hydroxide ligands resulted in few examples characterized by crystallography: (a) Wiegardt, K.; Bossek, U.; Nuber, B.; Weiss, J.; Bonvoisin, J.; Corbella, M.; Vitols, S. E.; Girerd, J. J., *J. Am. Chem. Soc.* **1988**, *110*, 7398-7411; (b) Pal, S.; Armstrong, W. H., *Inorg. Chem.* **1992**, *31*, 5417-5423; (c) Pal, S.; Chan, M. K.; Armstrong, W. H., *J. Am. Chem. Soc.* **1992**, *114*, 6398-6406; (d) Chen, H. Y.; Collomb, M. N.; Duboc, C.; Blondin, G.; Riviere, E.; Faller, J. W.; Crabtree, R. H.; Brudvig, G. W., *Inorg. Chem.* **2005**, *44*, 9567-9573; (e) Corbella, M.; Costa, R.; Ribas, J.; Fries, P. H.; Latour, J. M.; Ohrstrom, L.; Solans, X.; Rodriguez, V., *Inorg. Chem.* **1996**, *35*, 1857-1865; (f) Ng, G. K. Y.; Ziller, J. W.; Borovik, A. S., *Chem. Commun.* **2012**, *48*, 2546-2548.
12. Tsui, E. Y.; Kanady, J. S.; Day, M. W.; Agapie, T., *Chem. Commun.* **2011**, *47*, 4189-4191.
13. (a) de Ruiter, G.; Thompson, N. B.; Lionetti, D.; Agapie, T., *J. Am. Chem. Soc.* **2015**, *137*, 14094-14106; (b) de Ruiter, G.; Thompson, N. B.; Takase, M. K.; Agapie, T., *J. Am. Chem. Soc.* **2016**, *138*, 1486-1489; (c) de Ruiter, G.; Carsch, K. M.; Gul, S.; Chatterjee, R.; Thompson, N. B.; Takase, M. K.; Yano, J.; Agapie, T., *Angew. Chem. Int. Ed.* **2017**, *56*, 4772-4776. (d) Kanady, J. S.; Tran, R.; Stull, J. A.; Lu, L.; Stich, T. A.; Day, M. W.; Yano, J.; Britt, R. D.; Agapie, T., *Chem. Sci.* **2013**, *4*, 3986-3996.
14. Sickerman, N. S.; Park, Y. J.; Ng, G. K. Y.; Bates, J. E.; Hilkert, M.; Ziller, J. W.; Furche, F.; Borovik, A. S., *Dalton Trans.* **2012**, *41*, 4358-4364.
15. Zhou, M. J.; Diwu, Z. J.; PanchukVoloshina, N.; Haugland, R. P., *Anal. Biochem.* **1997**, *253*, 162-168.
16. Pegis, M. L.; Roberts, J. A. S.; Wasylenko, D. J.; Mader, E. A.; Appel, A. M.; Mayer, J. M., *Inorg. Chem.* **2015**, *54*, 11883-11888.
17. A MnO<sub>2</sub> catalyst has been shown to produce H<sub>2</sub>O<sub>2</sub> only at basic pH: Izgorodin, A.; Izgorodina, E.; MacFarlane, D. R., *Energ. Environ. Sci.* **2012**, *5*, 9496-9501.

TOC graphic:

